

Exercise 7(B)

Solution:

(i) Let's assume the fourth proportional to 1.5, 4.5 and 3.5 be x.

$$1.5: 4.5 = 3.5: x$$

$$1.5 \times x = 3.5 \times 4.5$$

$$x = (3.5 \times 4.5) / 1.5$$

$$x = 10.5$$

(ii) Let's assume the fourth proportional to $3a$, $6a^2$ and $2ab^2$ be x.

$$3a: 6a^2 = 2ab^2: x$$

$$3a \times x = 2ab^2 \times 6a^2$$

$$3a \times x = 12a^3b^2$$

$$x = 4a^2b^2$$

(i) **Solution:**

(i) Let's take the third proportional to $2\frac{2}{3}$ and 4 be x.

So, $2\frac{2}{3}$, 4, x are in continued proportion.

$$8/3: 4 = 4: x$$

$$(8/3)/ 4 = 4/x$$

$$x = 16 \times 3/8 = 6$$

(ii) Let's take the third proportional to $a - b$ and $a^2 - b^2$ be x.

So, $a - b$, $a^2 - b^2$, x are in continued proportion.

$$a - b: a^2 - b^2 = a^2 - b^2: x$$

$$\frac{a - b}{a^2 - b^2} = \frac{a^2 - b^2}{x}$$

$$x = \frac{(a^2 - b^2)^2}{a - b}$$

$$x = \frac{(a + b)(a - b)(a^2 - b^2)}{a - b}$$

$$x = (a + b)(a^2 - b^2)$$

2. Find the mean proportional between:

(i) $6 + 3\sqrt{3}$ and $8 - 4\sqrt{3}$

(ii) $a - b$ and $a^3 - a^2b$

Solution:

(i) Let the mean proportional between $6 + 3\sqrt{3}$ and $8 - 4\sqrt{3}$ be x .

So, $6 + 3\sqrt{3}$, x and $8 - 4\sqrt{3}$ are in continued proportion.

$$6 + 3\sqrt{3} : x = x : 8 - 4\sqrt{3}$$

$$x \times x = (6 + 3\sqrt{3})(8 - 4\sqrt{3})$$

$$x^2 = 48 + 24\sqrt{3} - 24\sqrt{3} - 36$$

$$x^2 = 12$$

$$x = 2\sqrt{3}$$

(ii) Let the mean proportional between $a - b$ and $a^3 - a^2b$ be x .

$a - b$, x , $a^3 - a^2b$ are in continued proportion.

$$a - b : x = x : a^3 - a^2b$$

$$x \times x = (a - b)(a^3 - a^2b)$$

$$x^2 = (a - b)a^2(a - b) = [a(a - b)]^2$$

$$x = a(a - b)$$

3. If $x + 5$ is the mean proportional between $x + 2$ and $x + 9$; find the value of x .

Solution:

Given, $x + 5$ is the mean proportional between $x + 2$ and $x + 9$.

So, $(x + 2)$, $(x + 5)$ and $(x + 9)$ are in continued proportion.

$$(x + 2) : (x + 5) = (x + 5) : (x + 9)$$

$$(x + 2) / (x + 5) = (x + 5) / (x + 9)$$

$$(x + 5)^2 = (x + 2)(x + 9)$$

$$x^2 + 25 + 10x = x^2 + 2x + 9x + 18$$

$$25 - 18 = 11x - 10x$$

$$x = 7$$

4. If x^2 , 4 and 9 are in continued proportion, find x .

Solution:

Given, x^2 , 4 and 9 are in continued proportion

So, we have

$$x^2/4 = 4/9$$

$$x^2 = 16/9$$

$$\text{Thus, } x = 4/3$$

5. What least number must be added to each of the numbers 6, 15, 20 and 43 to make them proportional?

Solution:

Let assume the number added to be x .

So, $(6 + x) : (15 + x) :: (20 + x) : (43 + x)$

$$(6 + x) / (15 + x) = (20 + x) / (43 + x)$$

$$(6 + x)(43 + x) = (20 + x)(43 + x)$$

$$258 + 6x + 43x + x^2 = 300 + 20x = 15x + x^2$$

$$49x - 35x = 300 - 258$$

$$14x = 42$$

$$x = 3$$

Therefore, the required number which should be added is 3.

6. (i) If a, b, c are in continued proportion,

Show that:
$$\frac{a^2 + b^2}{b(a+c)} = \frac{b(a+c)}{b^2 + c^2}$$

Solution:

Given,

a, b, c are in continued proportion.

So, we have

$$a/b = b/c$$

$$\Rightarrow b^2 = ac$$

Now,

$$(a^2 + b^2)(b^2 + c^2) = (a^2 + ac)(ac + c^2) \quad [\text{As } b^2 = ac]$$

$$= a(a+c)c(a+c)$$

$$= ac(a+c)^2$$

$$= b^2(a+c)^2$$

$$(a^2 + b^2)(b^2 + c^2) = [b(a+c)][b(a+c)]$$

Thus, L.H.S = R.H.S

$$\frac{a^2 + b^2}{b(a+c)} = \frac{b(a+c)}{b^2 + c^2}$$

- Hence Proved

(ii) If a, b, c are in continued proportion and a(b - c) = 2b, prove that: a - c = 2(a + b)/ a

Solution:

Given,

a, b, c are in continued proportion.

So, we have

$$a/b = b/c$$

$$\Rightarrow b^2 = ac$$

And, given $a(b - c) = 2b$

$$ab - ac = 2b$$

$$ab - b^2 = 2b$$

$$ab = 2b + b^2$$

$$ab = b(2 + b)$$

$$a = b + 2$$

$$a - b = 2$$

Now, taking the L.H.S we have

$$\text{L.H.S} = a - c$$

$$= a(a - c)/ a \quad [\text{Multiply and divide by } a]$$

$$\begin{aligned}
 &= a^2 - ac/a \\
 &= a^2 - b^2/a \\
 &= (a - b)(a + b)/a \\
 &= 2(a + b)/a \\
 &= \text{R.H.S}
 \end{aligned}$$

- Hence Proved

(iii) If $a/b = c/d$, show that: $\frac{a^3c + ac^3}{b^3d + bd^3} = \frac{(a + c)^4}{(b + d)^4}$

Solution:

Let's take $a/b = c/d = k$

So, $a = bk$ and $c = dk$

Taking L.H.S,

$$\begin{aligned}
 \text{L.H.S.} &= \frac{a^3c + ac^3}{b^3d + bd^3} = \frac{ac(a^2 + c^2)}{bd(b^2 + d^2)} \\
 &= \frac{(bk \times dk)(b^2k^2 + d^2k^2)}{bd(b^2 + d^2)} \\
 &= \frac{k^2 \times k^2(b^2 + d^2)}{(b^2 + d^2)} = k^4
 \end{aligned}$$

Now, taking the R.H.S

$$\text{R.H.S.} = \frac{(a + c)^4}{(b + d)^4} = \frac{(bk + dk)^4}{(b + d)^4} = \left[\frac{k(b + d)}{b + d} \right]^4 = k^4$$

Thus, L.H.S = R.H.S

- Hence Proved

7. What least number must be subtracted from each of the numbers 7, 17 and 47 so that the remainders are in continued proportion?

Solution:

Let's assume the number subtracted to be x .

So, we have

$$(7 - x) : (17 - x) :: (17 - x) : (47 - x)$$

$$\frac{7 - x}{17 - x} = \frac{17 - x}{47 - x}$$

$$(7 - x)(17 - x) = (17 - x)^2$$

$$329 - 47x - 7x + x^2 = 289 - 34x + x^2$$

$$329 - 289 = -34x + 54x$$

$$20x = 40$$

$$x = 2$$

Therefore, the required number which must be subtracted is 2.