

$$= \int e^x(\sec x + \sec x \tan x) dx$$

$$= \int e^x \sec x dx + \int e^x \sec x \tan x dx$$

Tip - If $f_1(x)$ and $f_2(x)$ are two functions, then an integral of the form $\int f_1(x)f_2(x)dx$ can be INTEGRATED BY PARTS as

$$f_1(x) \int f_2(x)dx - \int \left\{ \frac{d}{dx} f_1(x) \int f_2(x)dx \right\} dx \text{ where } f_1(x) \text{ and } f_2(x) \text{ are the first and second functions respectively.}$$

Taking $f_1(x) = \sec x$ and $f_2(x) = e^x$ in the first integral and keeping the second integral intact,

$$\int e^x \sec x dx + \int e^x \sec x \tan x dx$$

$$= \sec x \int e^x dx - \int \left[\frac{d}{dx} (\sec x) \int e^x dx \right] dx + \int e^x \sec x \tan x dx$$

$$= e^x \sec x - \int e^x \sec x \tan x dx + \int e^x \sec x \tan x dx + c$$

$$= e^x \sec x + c, \text{ where } c \text{ is the integrating constant}$$

66. Question

Evaluate the following integrals:

$$\int e^x \left(\frac{2 - \sin 2x}{1 - \cos 2x} \right) dx$$

Answer

$$\int e^x \left(\frac{2 - \sin 2x}{1 - \cos 2x} \right) dx$$

$$= \int e^x \left(\frac{1 - \sin x \cos x}{\sin^2 x} \right) dx$$

$$= \int e^x (\operatorname{cosec}^2 x - \cot x) dx$$

$$= \int e^x \operatorname{cosec}^2 x dx - \int e^x \cot x dx$$

Tip - If $f_1(x)$ and $f_2(x)$ are two functions, then an integral of the form $\int f_1(x)f_2(x)dx$ can be INTEGRATED BY PARTS as

$$f_1(x) \int f_2(x)dx - \int \left\{ \frac{d}{dx} f_1(x) \int f_2(x)dx \right\} dx \text{ where } f_1(x) \text{ and } f_2(x) \text{ are the first and second functions respectively.}$$

Taking $f_1(x) = \cot x$ and $f_2(x) = e^x$ in the second integral and keeping the first integral intact,

$$\int e^x \operatorname{cosec}^2 x dx - \int e^x \cot x dx$$

$$= \int e^x \operatorname{cosec}^2 x dx - \cot x \int e^x dx + \int \left\{ \frac{d}{dx} (\cot x) \int e^x dx \right\} dx$$

$$= \int e^x \operatorname{cosec}^2 x dx - e^x \cot x - \int e^x \operatorname{cosec}^2 x dx$$

$$= -e^x \cot x + c, \text{ where } c \text{ is the integrating constant}$$

67. Question

Evaluate the following integrals:



$$\int e^x \left(\frac{1 + \sin x}{1 + \cos x} \right) dx$$

Answer

$$\left(\frac{1 + \sin x}{1 + \cos x} \right)$$

$$= \left(\frac{1 + \frac{2 \tan^{x/2}}{1 + \tan^2(x/2)}}{1 + \frac{1 - \tan^2(x/2)}{1 + \tan^2(x/2)}} \right)$$

$$= \frac{(1 + \tan^{x/2})^2}{2}$$

$$\therefore \int e^x \left(\frac{1 + \sin x}{1 + \cos x} \right) dx$$

$$= \int e^x \times \frac{(1 + \tan^{x/2})^2}{2}$$

$$= \int \frac{e^x (1 + \tan^2 x/2 + 2 \tan^{x/2})}{2} dx$$

$$= \int \frac{e^x (\sec^2 x/2 + 2 \tan^{x/2})}{2} dx$$

$$= \int \frac{e^x \sec^2 x/2 dx}{2} + \int e^x \tan^{x/2} dx$$



Tip - If $f_1(x)$ and $f_2(x)$ are two functions, then an integral of the form $\int f_1(x) f_2(x) dx$ can be INTEGRATED BY PARTS as

$$f_1(x) \int f_2(x) dx - \int \left\{ \frac{d}{dx} f_1(x) \int f_2(x) dx \right\} dx$$

where $f_1(x)$ and $f_2(x)$ are the first and second functions respectively.

Taking $f_1(x) = \tan(x/2)$ and $f_2(x) = e^x$ in the second integral and keeping the first integral intact,

$$\int \frac{e^x \sec^2 x/2 dx}{2} + \int e^x \tan^{x/2} dx$$

$$= \int \frac{e^x \sec^2 x/2 dx}{2} + \tan^{x/2} \int e^x dx - \int \left[\frac{d}{dx} (\tan^{x/2}) \int e^x dx \right] dx$$

$$= \int \frac{e^x \sec^2 x/2 dx}{2} + e^x \tan^{x/2} - \int \frac{e^x \sec^2 x/2 dx}{2} + c$$

$$= e^x \tan^{x/2} + c, \text{ where } c \text{ is the integrating constant}$$

68. Question

Evaluate the following integrals:

$$\int e^x \left(\frac{\sin 4x - 4}{1 - \cos 4x} \right) dx$$

Answer

$$\int e^x \left(\frac{\sin 4x - 1}{1 - \cos 4x} \right) dx$$

$$\begin{aligned}
&= \int e^x \left(\frac{2\sin 2x \cos 2x - 4}{2\sin^2 2x} \right) dx \\
&= \int e^x (\cot 2x - 2\operatorname{cosec}^2 2x) dx \\
&= \int e^x \cot 2x dx - \int 2e^x \operatorname{cosec}^2 2x dx
\end{aligned}$$

Tip - If $f_1(x)$ and $f_2(x)$ are two functions, then an integral of the form $\int f_1(x)f_2(x)dx$ can be INTEGRATED BY PARTS as

$$f_1(x) \int f_2(x)dx - \int \left\{ \frac{d}{dx} f_1(x) \int f_2(x)dx \right\} dx \text{ where } f_1(x) \text{ and } f_2(x) \text{ are the first and second functions respectively.}$$

Taking $f_1(x) = \cot 2x$ and $f_2(x) = e^x$ in the first integral and keeping the second integral intact,

$$\begin{aligned}
&\int e^x \cot 2x dx - \int 2e^x \operatorname{cosec}^2 2x dx \\
&= \cot 2x \int e^x dx - \int \left\{ \frac{d}{dx} (\cot 2x) \int e^x dx \right\} dx - \int 2e^x \operatorname{cosec}^2 2x dx \\
&= e^x \cot 2x + \int 2e^x \operatorname{cosec}^2 2x dx - \int 2e^x \operatorname{cosec}^2 2x dx + c \\
&= e^x \cot 2x + c, \text{ where } c \text{ is the integrating constant}
\end{aligned}$$

69. Question

Evaluate the following integrals:

$$\int \frac{e^x [\sqrt{1-x^2} \sin^{-1} x + 1]}{\sqrt{1-x^2}} dx$$



Answer

$$\begin{aligned}
&\int \frac{e^x [\sqrt{1-x^2} \sin^{-1} x + 1]}{\sqrt{1-x^2}} dx \\
&= \int e^x \left(\sin^{-1} x + \frac{1}{\sqrt{1-x^2}} \right) dx \\
&= \int e^x \sin^{-1} x dx + \int \frac{e^x}{\sqrt{1-x^2}} dx
\end{aligned}$$

Tip - If $f_1(x)$ and $f_2(x)$ are two functions, then an integral of the form $\int f_1(x)f_2(x)dx$ can be INTEGRATED BY PARTS as

$$f_1(x) \int f_2(x)dx - \int \left\{ \frac{d}{dx} f_1(x) \int f_2(x)dx \right\} dx \text{ where } f_1(x) \text{ and } f_2(x) \text{ are the first and second functions respectively.}$$

Taking $f_1(x) = \sin^{-1} x$ and $f_2(x) = e^x$ in the first integral and keeping the second integral intact,

$$\begin{aligned}
&\int e^x \sin^{-1} x dx + \int \frac{e^x}{\sqrt{1-x^2}} dx \\
&= \sin^{-1} x \int e^x dx - \int \left\{ \frac{d}{dx} (\sin^{-1} x) \int e^x dx \right\} dx + \int \frac{e^x}{\sqrt{1-x^2}} dx \\
&= e^x \sin^{-1} x - \int \frac{e^x}{\sqrt{1-x^2}} dx + \int \frac{e^x}{\sqrt{1-x^2}} dx + c \\
&= e^x \sin^{-1} x + c, \text{ where } c \text{ is the integrating constant}
\end{aligned}$$

70. Question

Evaluate the following integrals:

$$\int e^x \left(\frac{1+x \log x}{x} \right) dx$$

Answer

$$\begin{aligned} & \int e^x \left(\frac{1+x \log x}{x} \right) dx \\ &= \int e^x \left(\frac{1}{x} + \log x \right) dx \\ &= \int \frac{e^x}{x} dx + \int e^x \log x dx \end{aligned}$$

Tip - If $f_1(x)$ and $f_2(x)$ are two functions, then an integral of the form $\int f_1(x)f_2(x)dx$ can be INTEGRATED BY PARTS as

$$f_1(x) \int f_2(x)dx - \int \left\{ \frac{d}{dx} f_1(x) \int f_2(x)dx \right\} dx \text{ where } f_1(x) \text{ and } f_2(x) \text{ are the first and second functions respectively.}$$

Taking $f_1(x) = \log x$ and $f_2(x) = e^x$ in the second integral and keeping the first integral intact,

$$\begin{aligned} & \int \frac{e^x}{x} dx + \int e^x \log x dx \\ &= \int \frac{e^x}{x} dx + \log x \int e^x dx - \int \left[\frac{d}{dx} (\log x) \int e^x dx \right] dx \\ &= \int \frac{e^x}{x} dx + e^x \log x - \int \frac{e^x}{x} dx + c \\ &= e^x \log x + c, \text{ where } c \text{ is the integrating constant} \end{aligned}$$

71. Question

Evaluate the following integrals:

$$\int e^x \cdot \frac{x}{(1+x)^2} dx$$

Answer

$$\frac{x}{(1+x)^2} = \frac{A}{(1+x)} + \frac{B}{(1+x)^2}$$

$$\Rightarrow x = A(1+x) + B$$

$$\text{For } x=-1, \text{ equation: } -1 = B \text{ i.e. } B = -1$$

$$\text{For } x=0, \text{ equation: } 0 = A-1 \text{ i.e. } A = 1$$

$$\therefore \frac{x}{(1+x)^2}$$

$$= \frac{1}{(1+x)} - \frac{1}{(1+x)^2}$$

The given equation becomes

$$\begin{aligned} & \int e^x \left[\frac{1}{(1+x)} - \frac{1}{(1+x)^2} \right] dx \\ &= \int e^x \times \frac{1}{(1+x)} dx - \int e^x \times \frac{1}{(1+x)^2} dx \end{aligned}$$

Tip - If $f_1(x)$ and $f_2(x)$ are two functions, then an integral of the form $\int f_1(x)f_2(x)dx$ can be INTEGRATED BY PARTS as

$$f_1(x) \int f_2(x)dx - \int \left\{ \frac{d}{dx} f_1(x) \int f_2(x)dx \right\} dx \text{ where } f_1(x) \text{ and } f_2(x) \text{ are the first and second functions respectively.}$$

Taking $f_1(x) = 1/(1+x)$ and $f_2(x) = e^x$ in the first integral and keeping the second integral intact,

$$\begin{aligned} & \int \frac{e^x}{(1+x)} dx - \int \frac{e^x}{(1+x)^2} dx \\ &= \frac{1}{(1+x)} \int e^x dx - \int \left[\frac{d}{dx} \left(\frac{1}{1+x} \right) \int e^x dx \right] dx - \int \frac{e^x}{(1+x)^2} dx \\ &= \frac{e^x}{(1+x)} + \int \frac{e^x}{(1+x)^2} dx - \int \frac{e^x}{(1+x)^2} dx + c \\ &= \frac{e^x}{(1+x)} + c, \text{ where } c \text{ is the integrating constant} \end{aligned}$$

72. Question

Evaluate the following integrals:

$$\int e^x \frac{(x-1)}{(x+1)^3} dx$$

Answer

$$\frac{x-1}{(x+1)^3} = \frac{A}{(x+1)} + \frac{B}{(x+1)^2} + \frac{C}{(x+1)^3}$$

$$\Rightarrow x-1 = A(x+1)^2 + B(x+1) + C$$

$$\text{For } x=-1, \text{ equation: } -2 = C \text{ i.e. } C = -2$$

$$\text{For } x=0, \text{ equation: } -1 = A+B-2 \text{ i.e. } A+B = 1$$

$$\text{For } x=1, \text{ equation: } 0 = 4A+2B-2$$

$$\text{i.e. } 2(A+B+A) = 2$$

$$\Rightarrow 1+A = 1$$

$$\Rightarrow A = 0$$

$$\text{And, } B = 1$$

$$\therefore \frac{x-1}{(x+1)^3}$$

$$= \frac{1}{(x+1)^2} - \frac{2}{(x+1)^3}$$

The given equation becomes

$$\begin{aligned} & \int e^x \left[\frac{1}{(x+1)^2} - \frac{2}{(x+1)^3} \right] dx \\ &= \int e^x \times \frac{1}{(x+1)^2} dx - \int e^x \times \frac{2}{(x+1)^3} dx \end{aligned}$$

Tip - If $f_1(x)$ and $f_2(x)$ are two functions, then an integral of the form $\int f_1(x)f_2(x)dx$ can be INTEGRATED BY PARTS as

$$f_1(x) \int f_2(x)dx - \int \left\{ \frac{d}{dx} f_1(x) \int f_2(x)dx \right\} dx \text{ where } f_1(x) \text{ and } f_2(x) \text{ are the first and second functions respectively.}$$



Taking $f_1(x) = 1/(1+x)^2$ and $f_2(x) = e^x$ in the first integral and keeping the second integral intact,

$$\begin{aligned} & \int \frac{e^x}{(x+1)^2} dx - \int \frac{2e^x}{(x+1)^3} dx \\ &= \frac{1}{(x+1)^2} \int e^x dx - \int \left[\frac{d}{dx} \left(\frac{1}{(x+1)^2} \right) \int e^x dx \right] dx - \int \frac{2e^x}{(x+1)^3} dx \\ &= \frac{e^x}{(x+1)^2} + \int \frac{2e^x}{(x+1)^3} dx - \int \frac{2e^x}{(x+1)^3} dx + c \\ &= \frac{e^x}{(x+1)^2} + c, \text{ where } c \text{ is the integrating constant} \end{aligned}$$

73. Question

Evaluate the following integrals:

$$\int e^x \frac{(2-x)}{(1-x)^2} dx$$

Answer

$$\frac{2-x}{(1-x)^2} = \frac{A}{(1-x)} + \frac{B}{(1-x)^2}$$

$$\Rightarrow 2-x = A(1-x) + B$$

For $x=1$, equation: $1 = B$ i.e. $B = 1$

For $x=2$, equation: $0 = -A+1$ i.e. $A = 1$

$$\therefore \frac{2-x}{(1-x)^2}$$

$$= \frac{1}{(1-x)} + \frac{1}{(1-x)^2}$$



The given equation becomes

$$\begin{aligned} & \int e^x \left[\frac{1}{(1-x)} + \frac{1}{(1-x)^2} \right] dx \\ &= \int e^x \times \frac{1}{(1-x)^2} dx + \int e^x \times \frac{1}{1-x} dx \end{aligned}$$

Tip - If $f_1(x)$ and $f_2(x)$ are two functions, then an integral of the form $\int f_1(x)f_2(x)dx$ can be INTEGRATED BY PARTS as

$f_1(x) \int f_2(x)dx - \int \left\{ \frac{d}{dx} f_1(x) \int f_2(x)dx \right\} dx$ where $f_1(x)$ and $f_2(x)$ are the first and second functions respectively.

Taking $f_1(x) = 1/(1-x)$ and $f_2(x) = e^x$ in the second integral and keeping the first integral intact,

$$\begin{aligned} & \int \frac{e^x}{(1-x)^2} dx + \int \frac{e^x}{1-x} dx \\ &= \int \frac{e^x}{(1-x)^2} dx + \frac{1}{1-x} \int e^x dx - \int \left[\frac{d}{dx} \left(\frac{1}{1-x} \right) \int e^x dx \right] dx \\ &= \int \frac{e^x}{(1-x)^2} dx + \frac{e^x}{1-x} - \int \frac{e^x}{(1-x)^2} dx + c \\ &= \frac{e^x}{1-x} + c, \text{ where } c \text{ is the integrating constant} \end{aligned}$$

74. Question

Evaluate the following integrals:

$$\int e^x \cdot \frac{(x-3)}{(x-1)^3} dx$$

Answer

$$\frac{x-3}{(x-1)^3} = \frac{A}{(x-1)} + \frac{B}{(x-1)^2} + \frac{C}{(x-1)^3}$$

$$\Rightarrow x-3 = A(x-1)^2 + B(x-1) + C$$

For $x=1$, equation: $-2 = C$ i.e. $C = -2$

For $x=0$, equation: $-3 = A-B-2$ i.e. $B = A+1$

For $x=3$, equation: $0 = 4A+2B-2$

$$\text{i.e. } 2(A+B+A) = 2$$

$$\Rightarrow 1+3A = 1$$

$$\Rightarrow A = 0$$

And, $B = 1$

$$\therefore \frac{x-3}{(x-1)^3}$$

$$= \frac{1}{(x-1)^2} - \frac{2}{(x-1)^3}$$

The given equation becomes

$$\int e^x \left[\frac{1}{(x-1)^2} - \frac{2}{(x-1)^3} \right] dx$$

$$= \int e^x \times \frac{1}{(x-1)^2} dx - \int e^x \times \frac{2}{(x-1)^3} dx$$



Tip - If $f_1(x)$ and $f_2(x)$ are two functions, then an integral of the form $\int f_1(x)f_2(x)dx$ can be INTEGRATED BY PARTS as

$f_1(x) \int f_2(x)dx - \int \left\{ \frac{d}{dx} f_1(x) \int f_2(x)dx \right\} dx$ where $f_1(x)$ and $f_2(x)$ are the first and second functions respectively.

Taking $f_1(x) = 1/(1-x)^2$ and $f_2(x) = e^x$ in the first integral and keeping the second integral intact,

$$\int \frac{e^x}{(x-1)^2} dx - \int \frac{2e^x}{(x-1)^3} dx$$

$$= \frac{1}{(x-1)^2} \int e^x dx - \int \left[\frac{d}{dx} \left(\frac{1}{(x-1)^2} \right) \int e^x dx \right] dx - \int \frac{2e^x}{(x-1)^3} dx$$

$$= \frac{e^x}{(x-1)^2} + \int \frac{2e^x}{(x-1)^3} dx - \int \frac{2e^x}{(x-1)^3} dx + c$$

$$= \frac{e^x}{(x-1)^2} + c, \text{ where } c \text{ is the integrating constant}$$

75. Question

Evaluate the following integrals:

$$\int e^{3x} \left(\frac{3x-1}{9x^2} \right) dx$$

Answer

$$\int e^{3x} \left(\frac{3x-1}{9x^2} \right) dx$$

$$= \int \frac{e^{3x}}{3x} dx - \int \frac{e^{3x}}{9x^2} dx$$

Tip - If $f_1(x)$ and $f_2(x)$ are two functions, then an integral of the form $\int f_1(x)f_2(x)dx$ can be INTEGRATED BY PARTS as

$$f_1(x) \int f_2(x)dx - \int \left\{ \frac{d}{dx} f_1(x) \int f_2(x)dx \right\} dx \text{ where } f_1(x) \text{ and } f_2(x) \text{ are the first and second functions respectively.}$$

Taking $f_1(x) = 1/3x$ and $f_2(x) = e^{3x}$ in the first integral and keeping the second integral intact,

$$\int \frac{e^{3x}}{3x} dx - \int \frac{e^{3x}}{9x^2} dx$$

$$= \frac{1}{3x} \int e^{3x} dx - \int \left[\frac{d}{dx} \left(\frac{1}{3x} \right) \int e^{3x} dx \right] dx - \int \frac{e^{3x}}{9x^2} dx$$

$$= \frac{e^{3x}}{9x} + \int \frac{e^{3x}}{9x^2} dx - \int \frac{e^{3x}}{9x^2} dx + c$$

$$= \frac{e^{3x}}{9x} + c, \text{ where } c \text{ is the integrating constant}$$

76. Question

Evaluate the following integrals:

$$\int \frac{(x+1)}{(x+2)^2} e^x dx$$

**Answer**

$$\frac{x+1}{(x+2)^2} = \frac{A}{x+2} + \frac{B}{(x+2)^2}$$

$$\Rightarrow x+1 = A(x+2) + B$$

For $x=-2$, equation: $-1 = B$ i.e. $B = -1$

For $x=-1$, equation: $0 = A-1$ i.e. $A = 1$

$$\therefore \frac{x+1}{(x+2)^2}$$

$$= \frac{1}{x+2} - \frac{1}{(x+2)^2}$$

The given equation becomes

$$\int e^x \left[\frac{1}{x+2} - \frac{1}{(x+2)^2} \right] dx$$

$$= \int e^x \times \frac{1}{x+2} dx - \int e^x \times \frac{1}{(x+2)^2} dx$$

Tip - If $f_1(x)$ and $f_2(x)$ are two functions, then an integral of the form $\int f_1(x)f_2(x)dx$ can be INTEGRATED BY PARTS as

$$f_1(x) \int f_2(x)dx - \int \left\{ \frac{d}{dx} f_1(x) \int f_2(x)dx \right\} dx \text{ where } f_1(x) \text{ and } f_2(x) \text{ are the first and second functions respectively.}$$

Taking $f_1(x) = 1/(x+2)$ and $f_2(x) = e^x$ in the second integral and keeping the first integral intact,

$$\begin{aligned} & \int \frac{e^x}{x+2} dx - \int \frac{e^x}{(x+2)^2} dx \\ &= \frac{1}{x+2} \int e^x dx - \int \left[\frac{d}{dx} \left(\frac{1}{x+2} \right) \int e^x dx \right] dx - \int \frac{e^x}{(x+2)^2} dx \\ &= \frac{e^x}{x+2} + \int \frac{e^x}{(x+2)^2} dx - \int \frac{e^x}{(x+2)^2} dx + c \\ &= \frac{e^x}{x+2} + c, \text{ where } c \text{ is the integrating constant} \end{aligned}$$

77. Question

Evaluate the following integrals:

$$\int \frac{x e^{2x}}{(1+2x)^2} dx$$

Answer

$$\frac{x}{(1+2x)^2} = \frac{A}{1+2x} + \frac{B}{(1+2x)^2}$$

$$\Rightarrow x = A(1+2x) + B$$

For $x = -1/2$, equation: $-1/2 = B$ i.e. $B = -1/2$

For $x = 0$, equation: $0 = A - 1/2$ i.e. $A = 1/2$

$$\therefore \frac{x}{(1+2x)^2}$$

$$= \frac{1}{2(1+2x)} - \frac{1}{2(1+2x)^2}$$



The given equation becomes

$$\begin{aligned} & \int e^{2x} \left[\frac{1}{2(1+2x)} - \frac{1}{2(1+2x)^2} \right] dx \\ &= \int e^{2x} \times \frac{1}{2(1+2x)} dx - \int e^{2x} \times \frac{1}{2(1+2x)^2} dx \end{aligned}$$

Tip - If $f_1(x)$ and $f_2(x)$ are two functions, then an integral of the form $\int f_1(x)f_2(x)dx$ can be INTEGRATED BY PARTS as

$$f_1(x) \int f_2(x)dx - \int \left\{ \frac{d}{dx} f_1(x) \int f_2(x)dx \right\} dx \text{ where } f_1(x) \text{ and } f_2(x) \text{ are the first and second functions respectively.}$$

Taking $f_1(x) = 1/(1+2x)$ and $f_2(x) = e^{2x}$ in the second integral and keeping the first integral intact,

$$\begin{aligned} & \int e^{2x} \times \frac{1}{2(1+2x)} dx - \int e^{2x} \times \frac{1}{2(1+2x)^2} dx \\ &= \frac{1}{2} \left[\frac{1}{1+2x} \int e^{2x} dx - \int \left[\frac{d}{dx} \left(\frac{1}{1+2x} \right) \int e^{2x} dx \right] dx - \int \frac{e^{2x}}{(1+2x)^2} dx \right] \\ &= \frac{1}{2} \left[\frac{e^{2x}}{2(2x+1)} + \int \frac{e^{2x}}{(2x+1)^2} dx - \int \frac{e^{2x}}{(2x+1)^2} dx \right] \\ &= \frac{e^{2x}}{4(2x+1)} + c, \text{ where } c \text{ is the integrating constant} \end{aligned}$$

78. Question

Evaluate the following integrals:

$$\int e^{2x} \left(\frac{2x-1}{4x^2} \right) dx$$

Answer

$$\begin{aligned} & \int e^{2x} \left(\frac{2x-1}{4x^2} \right) dx \\ &= \int \frac{e^{2x}}{2x} dx - \int \frac{e^{2x}}{4x^2} dx \end{aligned}$$

Tip - If $f_1(x)$ and $f_2(x)$ are two functions, then an integral of the form $\int f_1(x)f_2(x)dx$ can be INTEGRATED BY PARTS as

$$f_1(x) \int f_2(x)dx - \int \left\{ \frac{d}{dx} f_1(x) \int f_2(x)dx \right\} dx \text{ where } f_1(x) \text{ and } f_2(x) \text{ are the first and second functions respectively.}$$

Taking $f_1(x) = 1/2x$ and $f_2(x) = e^{2x}$ in the first integral and keeping the second integral intact,

$$\begin{aligned} & \int \frac{e^{2x}}{2x} dx - \int \frac{e^{2x}}{4x^2} dx \\ &= \frac{1}{2x} \int e^{2x} dx - \int \left[\frac{d}{dx} \left(\frac{1}{2x} \right) \int e^{2x} dx \right] dx - \int \frac{e^{2x}}{4x^2} dx \\ &= \frac{e^{2x}}{4x} + \int \frac{e^{2x}}{4x^2} dx - \int \frac{e^{2x}}{4x^2} dx + c \\ &= \frac{e^{2x}}{4x} + c, \text{ where } c \text{ is the integrating constant} \end{aligned}$$

79. Question

Evaluate the following integrals:

$$\int e^x \left(\log x + \frac{1}{x^2} \right) dx$$

Answer

$$\begin{aligned} & \int e^x \left(\log x + \frac{1}{x^2} \right) dx \\ &= \int e^x \log x dx - \int \frac{e^x}{x^2} dx \end{aligned}$$

Tip - If $f_1(x)$ and $f_2(x)$ are two functions, then an integral of the form $\int f_1(x)f_2(x)dx$ can be INTEGRATED BY PARTS as

$$f_1(x) \int f_2(x)dx - \int \left\{ \frac{d}{dx} f_1(x) \int f_2(x)dx \right\} dx \text{ where } f_1(x) \text{ and } f_2(x) \text{ are the first and second functions respectively.}$$

Taking $f_1(x) = \log x$ and $f_2(x) = e^x$ in the first integral and keeping the second integral intact,

$$\begin{aligned} & \int e^x \log x dx - \int \frac{e^x}{x^2} dx \\ &= \log x \int e^x dx - \int \left[\frac{d}{dx} (\log x) \int e^x dx \right] dx - \int \frac{e^x}{x^2} dx \\ &= e^x \log x - \int \frac{e^x}{x} dx - \int \frac{e^x}{x^2} dx \\ &= e^x \log x - \left[\frac{1}{x} \int e^x dx - \int \left[\frac{d}{dx} \left(\frac{1}{x} \right) \int e^x dx \right] dx \right] - \int \frac{e^x}{x^2} dx \end{aligned}$$



$$= e^x \log x - \frac{e^x}{x} + \int \frac{e^x}{x^2} dx - \int \frac{e^x}{x^2} dx + c$$

$$= e^x \left(\log x - \frac{1}{x} \right) + c, \text{ where } c \text{ is the integrating constant}$$

80. Question

Evaluate the following integrals:

$$\int \frac{\log x}{(1 + \log x)^2} dx$$

Answer

$$\frac{\log x}{(1 + \log x)^2} = \frac{A}{(1 + \log x)} + \frac{B}{(1 + \log x)^2}$$

$$\Rightarrow \log x = A(1 + \log x) + B$$

$$\text{For } x=1, \text{ equation: } 0 = A+B$$

$$\text{For } x=1/e, \text{ equation: } -1 = B \text{ i.e. } B = -1$$

$$\text{So, } A = 1$$

$$\therefore \frac{\log x}{(1 + \log x)^2}$$

$$= \frac{1}{(1 + \log x)} - \frac{1}{(1 + \log x)^2}$$

The given equation becomes

$$\int \left[\frac{1}{(1 + \log x)} - \frac{1}{(1 + \log x)^2} \right] dx$$

$$= \int \frac{1}{(1 + \log x)} dx - \int \frac{1}{(1 + \log x)^2} dx$$

Tip - If $f_1(x)$ and $f_2(x)$ are two functions, then an integral of the form $\int f_1(x)f_2(x)dx$ can be INTEGRATED BY PARTS as

$$f_1(x) \int f_2(x)dx - \int \left\{ \frac{d}{dx} f_1(x) \int f_2(x)dx \right\} dx \text{ where } f_1(x) \text{ and } f_2(x) \text{ are the first and second functions respectively.}$$

Taking $f_1(x) = 1/(1+\log x)$ and $f_2(x) = 1$ in the second integral and keeping the first integral intact,

$$\int \frac{1}{(1 + \log x)} dx - \int \frac{1}{(1 + \log x)^2} dx$$

$$= \frac{1}{(1 + \log x)} \int dx - \int \left[\frac{d}{dx} \left(\frac{1}{(1 + \log x)} \right) \int dx \right] dx - \int \frac{1}{(1 + \log x)^2} dx$$

$$= \frac{x}{(1 + \log x)} + \int \frac{1}{(1 + \log x)^2} dx - \int \frac{1}{(1 + \log x)^2} dx + c$$

$$= \frac{x}{(1 + \log x)} + c, \text{ where } c \text{ is the integrating constant}$$

81. Question

Evaluate the following integrals:

$$\int \{ \sin(\log x) + \cos(\log x) \} dx$$

Answer

Tip - If $f_1(x)$ and $f_2(x)$ are two functions, then an integral of the form $\int f_1(x)f_2(x)dx$ can be INTEGRATED BY PARTS as

$$f_1(x) \int f_2(x)dx - \int \left\{ \frac{d}{dx} f_1(x) \int f_2(x)dx \right\} dx \text{ where } f_1(x) \text{ and } f_2(x) \text{ are the first and second functions respectively.}$$

Taking $f_1(x) = \sin(\log x)$ and $f_2(x) = 1$ in the first integral and keeping the second integral intact,

$$\begin{aligned} & \int \sin(\log x)dx + \int \cos(\log x)dx \\ &= \sin(\log x) \int dx - \int \left[\frac{d}{dx} (\sin(\log x)) \int dx \right] dx + \int \cos(\log x)dx \\ &= x \sin(\log x) - \int \cos(\log x) dx + \int \cos(\log x)dx + c \\ &= e^{\log x} \sin(\log x) + c, \text{ where } c \text{ is the integrating constant} \end{aligned}$$

82. Question

Evaluate the following integrals:

$$\int \left\{ \frac{1}{\log x} - \frac{1}{(\log x)^2} \right\} dx$$

Answer

Tip - If $f_1(x)$ and $f_2(x)$ are two functions, then an integral of the form $\int f_1(x)f_2(x)dx$ can be INTEGRATED BY PARTS as

$$f_1(x) \int f_2(x)dx - \int \left\{ \frac{d}{dx} f_1(x) \int f_2(x)dx \right\} dx \text{ where } f_1(x) \text{ and } f_2(x) \text{ are the first and second functions respectively.}$$

Taking $f_1(x) = 1/(\log x)$ and $f_2(x) = 1$ in the first integral and keeping the second integral intact,

$$\begin{aligned} & \int \frac{1}{\log x} dx - \int \frac{1}{(\log x)^2} dx \\ &= \frac{1}{\log x} \int dx - \int \left[\frac{d}{dx} \left(\frac{1}{\log x} \right) \int dx \right] dx - \int \frac{1}{(\log x)^2} dx \\ &= \frac{x}{\log x} + \int \frac{1}{(\log x)^2} dx - \int \frac{1}{(\log x)^2} dx + c \\ &= \frac{x}{\log x} + c, \text{ where } c \text{ is the integrating constant} \end{aligned}$$

83. Question

Evaluate the following integrals:

$$\int \left\{ \log(\log x) + \frac{1}{(\log x)^2} \right\} dx$$

Answer

Tip - If $f_1(x)$ and $f_2(x)$ are two functions, then an integral of the form $\int f_1(x)f_2(x)dx$ can be INTEGRATED BY PARTS as

$$f_1(x) \int f_2(x)dx - \int \left\{ \frac{d}{dx} f_1(x) \int f_2(x)dx \right\} dx \text{ where } f_1(x) \text{ and } f_2(x) \text{ are the first and second functions respectively.}$$

Taking $f_1(x) = \log(\log x)$ and $f_2(x) = 1$ in the first integral and keeping the second integral intact,

$$\int \log(\log x)dx + \int \frac{1}{(\log x)^2} dx$$

$$\begin{aligned}
&= \log(\log x) \int dx - \int \left[\frac{d}{dx}(\log(\log x)) \int dx \right] dx + \int \frac{1}{(\log x)^2} dx \\
&= x \log(\log x) - \int \frac{1}{\log x} dx + \int \frac{1}{(\log x)^2} dx \\
&= x \log(\log x) - \left[\frac{1}{\log x} \int dx - \int \left[\frac{d}{dx} \left(\frac{1}{\log x} \right) \int dx \right] dx \right] + \int \frac{1}{(\log x)^2} dx \\
&= x \log(\log x) - \frac{x}{\log x} - \int \frac{1}{(\log x)^2} dx + \int \frac{1}{(\log x)^2} dx + c \\
&= x \left[\log(\log x) - \frac{1}{\log x} \right] + c, \text{ where } c \text{ is the integrating constant}
\end{aligned}$$

84. Question

Evaluate the following integrals:

$$\int \left(\frac{\sin^{-1} \sqrt{x} - \cos^{-1} \sqrt{x}}{\sin^{-1} \sqrt{x} + \cos^{-1} \sqrt{x}} \right) dx$$

Answer

It is known that $\sin^{-1} x + \cos^{-1} x = \pi/2$

$$\begin{aligned}
&\therefore \left(\frac{\sin^{-1} \sqrt{x} - \cos^{-1} \sqrt{x}}{\sin^{-1} \sqrt{x} + \cos^{-1} \sqrt{x}} \right) \\
&= \frac{2}{\pi} (\sin^{-1} \sqrt{x} - \cos^{-1} \sqrt{x})
\end{aligned}$$

Tip - If $f_1(x)$ and $f_2(x)$ are two functions, then an integral of the form $\int f_1(x) f_2(x) dx$ can be INTEGRATED BY PARTS as

$f_1(x) \int f_2(x) dx - \int \left\{ \frac{d}{dx} f_1(x) \int f_2(x) dx \right\} dx$ where $f_1(x)$ and $f_2(x)$ are the first and second functions respectively.

Now, for the first term,

Taking $f_1(x) = \sin^{-1} \sqrt{x}$ and $f_2(x) = 1$,

$$\begin{aligned}
&\therefore \int \sin^{-1} \sqrt{x} dx \\
&= \sin^{-1} \sqrt{x} \int dx - \int \left\{ \frac{d}{dx} (\sin^{-1} \sqrt{x}) \int dx \right\} dx \\
&= x \sin^{-1} \sqrt{x} - \int \frac{1}{2\sqrt{x}\sqrt{1-x}} \times x dx \\
&= x \sin^{-1} \sqrt{x} - \frac{1}{2} \int \frac{\sqrt{x}}{\sqrt{1-x}} dx
\end{aligned}$$

Taking $(1-x) = a^2$,

$-dx = 2a da$ i.e. $dx = -2a da$

Again, $x = 1 - a^2$

$$\begin{aligned}
&\therefore \frac{1}{2} \int \frac{\sqrt{x}}{\sqrt{1-x}} dx \\
&= \frac{1}{2} \int \frac{\sqrt{1-a^2}}{a} (-2a da)
\end{aligned}$$

$$= - \int \sqrt{1 - a^2} da$$

$$= - \left[\frac{1}{2} a \sqrt{1 - a^2} + \frac{1}{2} \sin^{-1} a \right]$$

Replacing the value of a, we get,

$$\therefore - \left[\frac{1}{2} a \sqrt{1 - a^2} + \frac{1}{2} \sin^{-1} a \right]$$

$$= - \left[\frac{1}{2} x \sqrt{1 - x} + \frac{1}{2} \sin^{-1} \sqrt{1 - x} \right] + c$$

The total integration yields as

$$= x \sin^{-1} \sqrt{x} + \left[\frac{1}{2} x \sqrt{1 - x} + \frac{1}{2} \sin^{-1} \sqrt{1 - x} \right] + c', \text{ where } c' \text{ is the integrating constant}$$

For the second term,

Taking $f_1(x) = \cos^{-1} \sqrt{x}$ and $f_2(x) = 1$,

$$\therefore \int \cos^{-1} \sqrt{x} dx$$

$$= \cos^{-1} \sqrt{x} \int dx - \int \left\{ \frac{d}{dx} (\cos^{-1} \sqrt{x}) \int dx \right\} dx$$

$$= x \cos^{-1} \sqrt{x} - \int \frac{-1}{2\sqrt{x}\sqrt{1-x}} \times x dx$$

$$= x \cos^{-1} \sqrt{x} + \frac{1}{2} \int \frac{\sqrt{x}}{\sqrt{1-x}} dx$$

Taking $(1-x)=a^2$,

$$-dx=2ada \text{ i.e. } dx=-2ada$$

Again, $x=1-a^2$

$$\therefore \frac{1}{2} \int \frac{\sqrt{x}}{\sqrt{1-x}} dx$$

$$= \frac{1}{2} \int \frac{\sqrt{1-a^2}}{a} (-2ada)$$

$$= - \int \sqrt{1 - a^2} da$$

$$= - \left[\frac{1}{2} a \sqrt{1 - a^2} + \frac{1}{2} \sin^{-1} a \right]$$

Replacing the value of a, we get,

$$\therefore - \left[\frac{1}{2} a \sqrt{1 - a^2} + \frac{1}{2} \sin^{-1} a \right]$$

$$= - \left[\frac{1}{2} x \sqrt{1 - x} + \frac{1}{2} \sin^{-1} \sqrt{1 - x} \right] + c$$

The total integration yields as

$$= x \cos^{-1} \sqrt{x} - \left[\frac{1}{2} x \sqrt{1 - x} + \frac{1}{2} \sin^{-1} \sqrt{1 - x} \right] + c'', \text{ where } c'' \text{ is the integrating constant}$$

$$\therefore \int \left(\frac{\sin^{-1} \sqrt{x} - \cos^{-1} \sqrt{x}}{\sin^{-1} \sqrt{x} + \cos^{-1} \sqrt{x}} \right) dx$$



$$\begin{aligned}
&= \frac{2}{\pi} \int (\sin^{-1} \sqrt{x} - \cos^{-1} \sqrt{x}) dx \\
&= \frac{2}{\pi} \left[x \sin^{-1} \sqrt{x} + \left[\frac{1}{2} x \sqrt{1-x} + \frac{1}{2} \sin^{-1} \sqrt{1-x} \right] - x \cos^{-1} \sqrt{x} \right. \\
&\quad \left. + \left[\frac{1}{2} x \sqrt{1-x} + \frac{1}{2} \sin^{-1} \sqrt{1-x} \right] \right] + c \\
&= \frac{2}{\pi} [\sqrt{x-x^2} + x(\sin^{-1} \sqrt{x} - \cos^{-1} \sqrt{x}) + \sin^{-1} \sqrt{1-x}] + c \text{ where } c \text{ is the integrating constant}
\end{aligned}$$

85. Question

Evaluate the following integrals:

$$\int 5^{5^x} \cdot 5^{5^x} \cdot 5^x dx$$

Answer

Tip - 5^x is to be replaced by a

$$\therefore 5^x = a$$

$$\Rightarrow 5^x \log 5 dx = da$$

$$\Rightarrow 5^x dx = \frac{da}{\log 5}$$

The equation becomes as follows:

$$\int 5^{5^a} \times 5^a \times \frac{1}{\log 5} da$$

Tip - 5^a is to be replaced by k

$$\therefore 5^a = k$$

$$\Rightarrow 5^a \log 5 da = dk$$

$$\Rightarrow 5^a da = \frac{dk}{\log 5}$$

The equation becomes as follows:

$$\int 5^k \times \frac{1}{(\log 5)^2} dk$$

$$= \frac{1}{(\log 5)^2} \int 5^k dk$$

$$= \frac{5^k}{(\log 5)^3} + c$$

Re-replacing the value of k,

$$\frac{5^{5^a}}{(\log 5)^3} + c$$

Re-replacing the value of a,

$$\frac{5^{5^{5^x}}}{(\log 5)^3} + c, \text{ where } c \text{ is the integrating constant}$$

86. Question

Evaluate the following integrals:



$$\int e^{2x} \left(\frac{1 + \sin 2x}{1 + \cos 2x} \right) dx$$

Answer

$$\left(\frac{1 + \sin 2x}{1 + \cos 2x} \right)$$

$$= \left(\frac{1 + \frac{2 \tan x}{1 + \tan^2 x}}{1 + \frac{1 - \tan^2 x}{1 + \tan^2 x}} \right)$$

$$= \frac{(1 + \tan x)^2}{2}$$

$$\therefore \int e^{2x} \left(\frac{1 + \sin 2x}{1 + \cos 2x} \right) dx$$

$$= \int e^{2x} \times \frac{(1 + \tan x)^2}{2}$$

$$= \int \frac{e^{2x}(1 + \tan^2 x + 2 \tan x)}{2} dx$$

$$= \int \frac{e^{2x}(\sec^2 x + 2 \tan x)}{2} dx$$

$$= \int \frac{e^{2x} \sec^2 x dx}{2} + \int e^{2x} \tan x dx$$

Tip - If $f_1(x)$ and $f_2(x)$ are two functions, then an integral of the form $\int f_1(x) f_2(x) dx$ can be INTEGRATED BY PARTS as

$f_1(x) \int f_2(x) dx - \int \left\{ \frac{d}{dx} f_1(x) \int f_2(x) dx \right\} dx$ where $f_1(x)$ and $f_2(x)$ are the first and second functions respectively.

Taking $f_1(x) = \tan x$ and $f_2(x) = e^{2x}$ in the second integral and keeping the first integral intact,

$$\int \frac{e^{2x} \sec^2 x dx}{2} + \int e^{2x} \tan x dx$$

$$= \int \frac{e^{2x} \sec^2 x dx}{2} + \tan x \int e^{2x} dx - \int \left[\frac{d}{dx} (\tan x) \int e^{2x} dx \right] dx$$

$$= \int \frac{e^{2x} \sec^2 x dx}{2} + \frac{1}{2} e^{2x} \tan x - \int \frac{e^{2x} \sec^2 x dx}{2} + c$$

$$= \frac{1}{2} e^x \tan x / 2 + c, \text{ where } c \text{ is the integrating constant}$$

87. Question

Evaluate the following integrals:

$$\int e^{2x} \left(\frac{1 - \sin 2x}{1 - \cos 2x} \right) dx$$

Answer

$$\left(\frac{1 - \sin 2x}{1 - \cos 2x} \right)$$

$$\begin{aligned}
&= \left(\frac{1 - \frac{2 \tan x}{1 + \tan^2 x}}{1 - \frac{1 - \tan^2 x}{1 + \tan^2 x}} \right) \\
&= \frac{(1 - \tan x)^2}{2} \\
\therefore \int e^{2x} \left(\frac{1 - \sin 2x}{1 - \cos 2x} \right) dx \\
&= \int e^{2x} \times \frac{(1 - \tan x)^2}{2} \\
&= \int \frac{e^{2x}(1 + \tan^2 x - 2 \tan x)}{2} dx \\
&= \int \frac{e^{2x}(\sec^2 x - 2 \tan x)}{2} dx \\
&= \int \frac{e^{2x} \sec^2 x dx}{2} - \int e^{2x} \tan x dx
\end{aligned}$$

Tip - If $f_1(x)$ and $f_2(x)$ are two functions, then an integral of the form $\int f_1(x)f_2(x)dx$ can be INTEGRATED BY PARTS as

$f_1(x) \int f_2(x)dx - \int \left\{ \frac{d}{dx} f_1(x) \int f_2(x)dx \right\} dx$ where $f_1(x)$ and $f_2(x)$ are the first and second functions respectively.

Taking $f_1(x) = \tan x$ and $f_2(x) = e^{2x}$ in the second integral and keeping the first integral intact,

$$\begin{aligned}
&\int \frac{e^{2x} \sec^2 x dx}{2} - \int e^{2x} \tan x dx \\
&= \int \frac{e^{2x} \sec^2 x dx}{2} - \tan x \int e^{2x} dx + \int \left[\frac{d}{dx} (\tan x) \int e^{2x} dx \right] dx \\
&= \int \frac{e^{2x} \sec^2 x dx}{2} - \frac{1}{2} e^{2x} \tan x + \int \frac{e^{2x} \sec^2 x dx}{2} + c \\
&= -\frac{1}{2} e^x \tan^x / 2 + c, \text{ where } c \text{ is the integrating constant}
\end{aligned}$$

Objective Questions II

1. Question

Mark (✓) against the correct answer in each of the following:

$$\int x e^x dx = ?$$

- A. $e^x (1 - x) + C$
- B. $e^x (x - 1) + C$
- C. $e^x (x - 1) + C$
- D. none of these

Answer

To find: Value of $\int x e^x dx$

Formula used:

$$(i) \int f(x)g(x)dx = f(x) \int g(x)dx - \int \left[f'(x) \int g(x)dx \right] dx$$

We have, $I = \int x e^x dx \dots (i)$

$$I = \int x e^x dx$$

$$\Rightarrow x \int e^x dx - \int \left[\frac{d(x)}{x} \int e^x dx \right] dx$$

$$\Rightarrow I = x e^x - \int 1 \cdot e^x dx$$

$$\Rightarrow I = x e^x - e^x + c$$

$$\therefore I = e^x (x-1) + c$$

Ans) $c e^x (x-1) + c$

2. Question

Mark (✓) against the correct answer in each of the following:

$$\int x e^{2x} dx = ?$$

A. $\frac{1}{2} x e^{2x} + \frac{1}{4} e^{2x} + C$

B. $\frac{1}{2} x e^{2x} - \frac{1}{4} e^{2x} + C$

C. $2x e^{2x} + 4e^{2x} + C$

D. none of these

Answer

To find: Value of $\int x e^{2x} dx$

Formula used:

$$(i) \int f(x)g(x)dx = f(x) \int g(x)dx - \int \left[f'(x) \int g(x)dx \right] dx$$

We have, $I = \int x e^{2x} dx \dots (i)$

$$I = \int x e^{2x} dx$$

$$\Rightarrow x \int e^{2x} dx - \int \left[\frac{d(x)}{x} \int e^{2x} dx \right] dx$$

$$\Rightarrow I = x \frac{e^{2x}}{2} - \int 1 \cdot \frac{e^{2x}}{2} dx$$

$$\Rightarrow I = x \frac{e^{2x}}{2} - \frac{1}{2} \int \frac{e^{2x}}{2} dx$$

$$\Rightarrow I = \frac{x}{2} e^{2x} - \frac{1}{2} \int e^{2x} dx$$

$$\Rightarrow I = \frac{x}{2} e^{2x} - \frac{1}{2} \frac{e^{2x}}{2} + c$$



$$\Rightarrow I = \frac{x}{2} e^{2x} - \frac{e^{2x}}{4} + c$$

$$\therefore I = \frac{x}{2} e^{2x} - \frac{e^{2x}}{4} + c$$

Ans) B $\frac{x}{2} e^{2x} - \frac{e^{2x}}{4} + c$

3. Question

Mark (✓) against the correct answer in each of the following:

$$\int x \cos 2x \, dx = ?$$

A. $\frac{1}{2} x \sin 2x + \frac{1}{4} \cos 2x + C$

B. $\frac{1}{2} x \sin 2x - \frac{1}{4} \cos 2x + C$

C. $2x \sin 2x + 4 \cos 2x + C$

D. none of these

Answer

To find: Value of $\int x \cos 2x \, dx$

Formula used:

$$(i) \int f(x)g(x)dx = f(x) \int g(x)dx - \int [f'(x) \int g(x)dx] dx$$

We have, $I = \int x \cos 2x \, dx$... (i)

Let $2x = t$

$$\Rightarrow x = \frac{t}{2}$$

$$\Rightarrow 2 = \frac{dt}{dx}$$

$$\Rightarrow dx = \frac{dt}{2}$$

$$I = \int \frac{t}{2} \cos t \frac{dt}{2}$$

$$I = \frac{1}{4} \int t \cos t \, dt$$

Taking 1st function as t and second function as $\cos t$

$$\Rightarrow I = \frac{1}{4} \left[t \int \cos t \, dt - \int \left(\frac{dt}{dt} \int \cos t \, dt \right) dt \right]$$

$$\Rightarrow I = \frac{1}{4} \left[t(\sin t) - \int (1(\sin t)) dt \right]$$

$$\Rightarrow I = \frac{1}{4} [t(\sin t) - (-\cos t)] + c$$

$$\Rightarrow I = \frac{1}{4} [t \sin t + \cos t] + c$$

$$\Rightarrow I = \frac{1}{4} [2x \sin 2x + \cos 2x] + c$$

$$\Rightarrow I = \frac{1}{2} x \sin 2x + \frac{1}{4} \cos 2x + c$$

Ans) A $\frac{1}{2} x \sin 2x + \frac{1}{4} \cos 2x + c$

4. Question

Mark (✓) against the correct answer in each of the following:

$$\int x \sec^2 x \, dx = ?$$

A. $x \tan x - \log |\cos x| + C$

B. $x \tan x + \log |\cos x| + C$

C. $x \tan x + \log |\sec x| + C$

D. none of these

Answer

To find: Value of $\int x \sec^2 x \, dx$

Formula used:

$$(i) \int f(x)g(x)dx = f(x) \int g(x)dx - \int \left[f'(x) \int g(x)dx \right] dx$$

We have, $I = \int x \sec^2 x \, dx \dots (i)$

Taking 1st function as x and second function as $\sec^2 x$

$$\Rightarrow I = \left[x \int \sec^2 x \, dx - \int \left(\frac{dx}{dx} \int \sec^2 x \, dx \right) dx \right]$$

$$\Rightarrow I = \left[x \tan x - \int (1 \tan x) dx \right]$$

$$\Rightarrow I = \left[x \tan x - \int \tan x dx \right]$$

$$\Rightarrow I = [x \tan x - (-\log |\cos x|)] + c$$

$$\Rightarrow I = x \tan x + \log |\cos x| + c$$

Ans) B $x \tan x + \log |\cos x| + c$

5. Question

Mark (✓) against the correct answer in each of the following:

$$\int x \sin 2x \, dx = ?$$

A. $\frac{1}{2} x \cos 2x + \frac{1}{4} \sin 2x + C$

B. $-\frac{1}{2} x \cos 2x - \frac{1}{4} \sin 2x + C$

$$C. -\frac{1}{2}x \cos 2x + \frac{1}{4} \sin 2x + C$$

D. none of these

Answer

To find: Value of $\int x \sin 2x dx$

Formula used:

$$(i) \int f(x)g(x)dx = f(x) \int g(x)dx - \int \left[f'(x) \int g(x)dx \right] dx$$

We have, $I = \int x \sin 2x dx \dots (i)$

Let $2x = t$

$$\Rightarrow x = \frac{t}{2}$$

$$\Rightarrow 2 = \frac{dt}{dx}$$

$$\Rightarrow dx = \frac{dt}{2}$$

$$I = \int \frac{t}{2} \sin t \frac{dt}{2}$$

$$I = \frac{1}{4} \int t \sin t dt$$

Taking 1st function as t and second function as $\sin t$

$$\Rightarrow I = \frac{1}{4} \left[t \int \sin t dt - \int \left(\frac{dt}{dt} \int \sin t dt \right) dt \right]$$

$$\Rightarrow I = \frac{1}{4} \left[t(-\cos t) - \int (1(-\cos t)) dt \right]$$

$$\Rightarrow I = \frac{1}{4} \left[-t \cos t - \int -\cos t dt \right]$$

$$\Rightarrow I = \frac{1}{4} [-t \cos t + \sin t] + c$$

$$\Rightarrow I = \frac{1}{4} [-2x \cos 2x + \sin 2x] + c$$

$$\Rightarrow I = -\frac{1}{2}x \cos 2x + \frac{1}{4} \sin 2x + c$$

Ans) C $-\frac{1}{2}x \cos 2x + \frac{1}{4} \sin 2x + c$

6. Question

Mark (✓) against the correct answer in each of the following:

$$\int x \log x dx = ?$$

A. $x \log x + \frac{1}{2}x^2 + C$



B. $\frac{1}{2}x^2 \log x + \frac{1}{4}x^2 + C$

C. $\frac{1}{2}x^2 \log x - \frac{1}{4}x^2 + C$

D. none of these

Answer

To find: Value of $\int x \log x \, dx$

Formula used:

$$(i) \int f(x)g(x)dx = f(x) \int g(x)dx - \int \left[f'(x) \int g(x)dx \right] dx$$

We have, $I = \int x \log x \, dx \dots (i)$

Taking 1st function as $\log x$ and second function as x

$$\Rightarrow I = \left[\log x \int x \, dx - \int \left(\frac{d \log x}{dx} \int x \, dx \right) dx \right]$$

$$\Rightarrow I = \left[\log x \frac{x^2}{2} - \int \left(\frac{1}{x} \frac{x^2}{2} \right) dx \right]$$

$$\Rightarrow I = \left[\log x \frac{x^2}{2} - \int \left(\frac{x}{2} \right) dx \right]$$

$$\Rightarrow I = \left[\log x \frac{x^2}{2} - \frac{1}{2} \int x \, dx \right]$$

$$\Rightarrow I = \left[\log x \frac{x^2}{2} - \frac{1}{2} \frac{x^2}{2} \right] + c$$

$$\Rightarrow I = \frac{1}{2}x^2 \log x - \frac{1}{4}x^2 + c$$

Ans) C $\frac{1}{2}x^2 \log x - \frac{1}{4}x^2 + c$

7. Question

Mark (✓) against the correct answer in each of the following:

$$\int x \operatorname{cosec}^2 x \, dx = ?$$

A. $x \cot x - \log |\sin x| + C$

B. $-\cot x + \log |\sin x| + C$

C. $x \tan x - \log |\sec x| + C$

D. none of these

Answer

To find: Value of $\int x \operatorname{cosec}^2 x \, dx$

Formula used:

$$(i) \int f(x)g(x)dx = f(x) \int g(x)dx - \int \left[f'(x) \int g(x)dx \right] dx$$

We have, $I = \int x \operatorname{cosec}^2 x dx \dots (i)$

$$I = \int x \operatorname{cosec}^2 x dx$$

$$\Rightarrow x \int \operatorname{cosec}^2 x dx - \int \left[\frac{d(x)}{x} \int \operatorname{cosec}^2 x dx \right] dx$$

$$\Rightarrow I = x(-\cot x) - \int 1.(-\cot x) dx$$

$$\Rightarrow I = -x(\cot x) + \log|\sin x| + c$$

Ans) D None of these

8. Question

Mark (✓) against the correct answer in each of the following:

$$\int x \sin x \cos x dx = ?$$

A. $-\frac{1}{4}x \sin 2x + \frac{1}{8} \cos 2x + C$

B. $\frac{1}{4}x \cos 2x - \frac{1}{8} \sin 2x + C$

C. $\frac{1}{2}x \sin 2x + \frac{1}{4} \cos 2x + C$

D. $-\frac{1}{4}x \cos 2x + \frac{1}{8} \sin 2x + C$



Answer

To find: Value of $\int x \sin x \cos x dx$

Formula used:

$$(i) \int f(x)g(x)dx = f(x) \int g(x)dx - \int \left[f'(x) \int g(x)dx \right] dx$$

We have, $I = \int x \sin x \cos x dx \dots (i)$

$$I = \frac{1}{2} \int x 2 \sin x \cos x dx$$

$$I = \frac{1}{2} \int x \sin 2x dx$$

$$\Rightarrow \frac{1}{2} \left[x \int \sin 2x dx - \int \left[\frac{d(x)}{x} \int \sin 2x dx \right] dx \right]$$

$$\Rightarrow \frac{1}{2} \left[\frac{-x \cos 2x}{2} - \int \left[1 \cdot \frac{-\cos 2x}{2} \right] dx \right]$$

$$\Rightarrow \frac{1}{2} \left[\frac{-x \cos 2x}{2} + \frac{\sin 2x}{4} \right] + c$$

$$\Rightarrow \frac{-x \cos 2x}{4} + \frac{\sin 2x}{8} + c$$

Ans) D $\frac{-x \cos 2x}{4} + \frac{\sin 2x}{8} + c$

9. Question

Mark (✓) against the correct answer in each of the following:

$\int x \cos^2 x \, dx = ?$

A. $\frac{x^2}{4} - \frac{x \sin 2x}{4} + \frac{\cos 2x}{8} + C$

B. $\frac{x^2}{4} + \frac{x \sin 2x}{4} + \frac{\cos 2x}{8} + C$

C. $\frac{x^2}{4} + \frac{x \sin 2x}{4} - \frac{\cos 2x}{8} + C$

D. none of these

Answer

To find: Value of $\int x \cos^2 x \, dx$

Formula used:

(i) $\int f(x)g(x)dx = f(x) \int g(x)dx - \int [f'(x) \int g(x)dx]dx$

We have, $I = \int x \cos^2 x \, dx \dots$ (i)

$I = \int x \frac{1}{2} (1 + \cos 2x) dx$

$I = \frac{1}{2} \int x \, dx + \frac{1}{2} \int x \cos 2x \, dx$

$I = \frac{1}{2} \frac{x^2}{2} + \frac{1}{2} \left[x \int \cos 2x \, dx - \int \left[\frac{d(x)}{dx} \int \cos 2x \, dx \right] dx \right]$

$I = \frac{1}{2} \frac{x^2}{2} + \frac{1}{2} \left[x \frac{\sin 2x}{2} - \int 1 \cdot \frac{\sin 2x}{2} \, dx \right]$

$I = \frac{1}{2} \frac{x^2}{2} + \frac{1}{2} \left[x \frac{\sin 2x}{2} - \frac{1}{2} \int \sin 2x \, dx \right]$

$I = \frac{1}{2} \frac{x^2}{2} + \frac{1}{2} \left[x \frac{\sin 2x}{2} - \frac{1}{2} \left(-\frac{\cos 2x}{2} \right) + c \right]$

$I = \frac{1}{2} \frac{x^2}{2} + \frac{1}{2} \left[\frac{x \sin 2x}{2} + \frac{\cos 2x}{4} + c \right]$

$I = \frac{x^2}{4} + \frac{x \sin 2x}{4} + \frac{\cos 2x}{8} + c$

Ans) D $\frac{x^2}{4} + \frac{x \sin 2x}{4} + \frac{\cos 2x}{8} + c$

10. Question

Mark (✓) against the correct answer in each of the following:



$$\int \frac{\log x}{x^2} dx = ?$$

A. $-\frac{1}{x}(\log x + 1) + C$

B. $\frac{1}{x}(\log x - 1) + C$

C. $\frac{1}{x}(\log x + 1) + C$

D. none of these

Answer

To find: Value of $\int \frac{\log x}{x^2} dx$

Formula used:

$$(i) \int f(x)g(x)dx = f(x) \int g(x)dx - \int [f'(x) \int g(x)dx] dx$$

We have, $I = \int \frac{\log x}{x^2} dx \dots (i)$

$$I = \int x^{-2} \log x dx$$

$$\Rightarrow \log x \int x^{-2} dx - \int \left[\frac{d(\log x)}{x} \int x^{-2} dx \right] dx$$

$$\Rightarrow \log x \frac{x^{-1}}{-1} - \int \left(\frac{1}{-x^2} \right) dx$$

$$\Rightarrow -\frac{\log x}{x} + \left(-\frac{1}{x} \right) + c$$

$$\Rightarrow -\frac{1}{x}(\log x + 1) + c$$

Ans) A - $\frac{1}{x}(\log x + 1) + c$

11. Question

Mark (✓) against the correct answer in each of the following:

$$\int \log x dx = ?$$

A. $\frac{1}{x} + C$

B. $\frac{1}{2}(\log x)^2 + C$

C. $x(\log x + 1) + C$

D. $x(\log x - 1) + C$

Answer



To find: Value of $\int \log x dx$

Formula used:

$$(i) \int f(x)g(x)dx = f(x) \int g(x)dx - \int \left[f'(x) \int g(x)dx \right] dx$$

We have, $I = \int \log x \cdot 1 \cdot dx \dots (i)$

Taking 1st function as $\log x$ and second function as 1

$$\Rightarrow I = \left[\log x \int 1 dx - \int \left(\frac{d \log x}{dx} \int 1 dx \right) dx \right]$$

$$\Rightarrow I = \left[\log x \cdot x - \int \left(\frac{1}{x} \int 1 dx \right) dx \right]$$

$$\Rightarrow I = \left[\log x \cdot x - \int \left(\frac{1}{x} \cdot x \right) dx \right]$$

$$\Rightarrow I = \left[\log x \cdot x - \int 1 dx \right]$$

$$\Rightarrow I = [\log x \cdot x - x] + c$$

$$\Rightarrow I = x(\log x - 1) + c$$

Ans) D $x(\log x - 1) + c$

12. Question

Mark (✓) against the correct answer in each of the following:

$$\int \log_{10} x dx = ?$$

A. $\frac{1}{x} \log_e 10 + C$

B. $\frac{1}{x} \log_{10} e + C$

C. $x (\log x - 1) \log_e 10 + C$

D. $x(\log x - 1) \log_{10} e + C$

Answer

To find: Value of $\int \log_{10} x dx$

Formula used: $\int \frac{1}{x} dx = \log|x| + c$

We have, $I = \int \log_{10} x dx \dots (i)$

$$I = \int \log_{10} x dx = \int \frac{\log x}{\log 10} dx$$

$$I = \frac{1}{\log_e 10} \int \log x \cdot 1 dx$$

Taking 1st function as $\log x$ and second function as 1



$$\Rightarrow I = \frac{1}{\log_e 10} \left[\log x \int 1 dx - \int \left(\frac{d \log x}{dx} \int 1 dx \right) dx \right]$$

$$\Rightarrow I = \frac{1}{\log_e 10} \left[\log x \cdot x - \int \left(\frac{1}{x} \int 1 dx \right) dx \right]$$

$$\Rightarrow I = \frac{1}{\log_e 10} \left[\log x \cdot x - \int \left(\frac{1}{x} x \right) dx \right]$$

$$\Rightarrow I = \frac{1}{\log_e 10} \left[\log x \cdot x - \int 1 dx \right]$$

$$\Rightarrow I = \frac{1}{\log_e 10} [\log x \cdot x - x] + c$$

$$\Rightarrow I = x(\log x - 1) \log_{10} e + c$$

Ans) D $x(\log x - 1) \log_{10} e + c$

13. Question

Mark (✓) against the correct answer in each of the following:

$$\int (\log x)^2 dx = ?$$

A. $\frac{2 \log x}{x} + C$

B. $\frac{1}{3} (\log x)^3 + C$

C. $x (\log x)^2 - 2x \log x + 2x + C$

D. $x (\log x)^2 + 2x \log x - 2x + C$



Answer

To find: Value of $\int (\log x)^2 dx$

Formula used:

$$(i) \int f(x)g(x)dx = f(x) \int g(x)dx - \int [f'(x) \int g(x)dx] dx$$

We have, $I = \int (\log x)^2 \cdot 1 \cdot dx \dots (i)$

Taking 1st function as $(\log x)^2$ and second function as 1

$$\Rightarrow I = \left[(\log x)^2 \int 1 dx - \int \left(\frac{d(\log x)^2}{dx} \int 1 dx \right) dx \right]$$

$$\Rightarrow I = \left[(\log x)^2 \int 1 dx - \int \left(\frac{2(\log x)}{x} \int 1 dx \right) dx \right]$$

$$\Rightarrow I = \left[(\log x)^2 \cdot x - 2 \int \log x dx \right]$$

$$\Rightarrow I = [(\log x)^2 \cdot x - 2(x \log x - x)] + c$$

$$\Rightarrow I = [(\log x)^2 \cdot x - 2x \log x + 2x] + c$$

$$\Rightarrow I = x(\log x)^2 - 2x \log x + 2x + c$$

Ans) C $x(\log x)^2 - 2x \log x + 2x + c$

14. Question

Mark (✓) against the correct answer in each of the following:

$$\int e^{\sqrt{x}} dx = ?$$

A. $e^{\sqrt{x}} + \sqrt{x} + C$

B. $\frac{1}{2} e^{\sqrt{x}} (\sqrt{x} + 1) + C$

C. $2e^{\sqrt{x}} (\sqrt{x} - 1) + C$

D. none of these

Answer

To find: Value of $\int e^{\sqrt{x}} dx$

Formula used: $\int \frac{1}{x} dx = \log|x| + c$

We have, $I = \int e^{\sqrt{x}} dx \dots (i)$

Putting $\sqrt{x} = t$

$$\Rightarrow \frac{1}{2\sqrt{x}} = \frac{dt}{dx}$$

$$\Rightarrow dx = 2\sqrt{x} dt$$

$$\Rightarrow dx = 2t dt$$

$$\Rightarrow I = 2 \int t \cdot e^t dt$$

$$\Rightarrow I = 2 \left[t \int e^t dt - \int \left[\frac{d(t)}{dt} \int e^t dt \right] dt \right]$$

$$\Rightarrow I = 2 \left[te^t - \int [1 \cdot e^t] dt \right]$$

$$\Rightarrow I = 2[te^t - e^t]$$

$$\Rightarrow I = e^t \cdot 2(t-1) + c$$

$$\therefore I = 2 e^{\sqrt{x}} (\sqrt{x} - 1) + c$$

Ans) C $2 e^{\sqrt{x}} (\sqrt{x} - 1) + c$

15. Question

Mark (✓) against the correct answer in each of the following:

$$\int \cos \sqrt{x} dx = ?$$

A. $\sin \sqrt{x} + \cos \sqrt{x} + C$



B. $\frac{1}{2}(\sqrt{x} \sin \sqrt{x} - \cos \sqrt{x}) + C$

C. $2[\sqrt{x} \sin \sqrt{x} + \cos \sqrt{x}] + C$

D. none of these

Answer

To find: Value of $\int \cos \sqrt{x} dx$

Formula used:

$$(i) \int f(x)g(x)dx = f(x) \int g(x)dx - \int [f'(x) \int g(x)dx]dx$$

We have, $I = \int \cos \sqrt{x} dx \dots (i)$

Putting $\sqrt{x}=t$

$$\Rightarrow \frac{1}{2\sqrt{x}} = \frac{dt}{dx}$$

$$\Rightarrow dx = 2\sqrt{x} dt$$

$$\Rightarrow dx = 2t dt$$

$$\Rightarrow I = \int \cos t \cdot 2t dt$$

$$\Rightarrow I = 2 \int t \cdot \cos t dt$$

$$\Rightarrow I = 2 \left[t \int \cos t dt - \int \left[\frac{d(t)}{dt} \int \cos t dt \right] dt \right]$$

$$\Rightarrow I = 2 \left[te^t - \int [1 e^t] dt \right]$$

$$\Rightarrow I = 2[te^t - e^t]$$

$$\Rightarrow I = e^t \cdot 2(t-1) + c$$

$$\therefore I = 2 e^{\sqrt{x}} (\sqrt{x}-1) + c$$

Ans) C $2 e^{\sqrt{x}} (\sqrt{x}-1) + c$

16. Question

Mark (✓) against the correct answer in each of the following:

$$\int \cos(\log x) dx = ?$$

A. $\frac{x}{2} [\cos(\log x) - \sin(\log x)] + C$

B. $\frac{x}{2} [\cos(\log x) + \sin(\log x)] + C$

C. $2x [\cos(\log x) + \sin(\log x)] + C$

D. $2x [\cos(\log x) - \sin(\log x)] + C$



Answer**To find:** Value of $\int \cos(\log x) dx$ **Formula used:**

$$(i) \int f(x)g(x)dx = f(x) \int g(x)dx - \int \left[f'(x) \int g(x)dx \right] dx$$

We have, $I = \int \cos(\log x) dx \dots (i)$

$$I = \int 1 \cdot \cos(\log x) dx$$

Taking $\cos(\log x)$ as first function and 1 as second function.

$$\Rightarrow I = \left[\cos \log x \int 1 dx - \int \left[\frac{d[\cos(\log x)]}{dx} \int 1 dx \right] dx \right]$$

$$\Rightarrow I = \left[x \cdot \cos(\log x) - \int \left[-\sin(\log x) \frac{1}{x} x \right] dx \right]$$

$$\Rightarrow I = \left[x \cdot \cos(\log x) + \int [\sin(\log x)] dx \right]$$

$$\Rightarrow I = \left[x \cdot \cos(\log x) + \int [1 \cdot \sin(\log x)] dx \right]$$

$$\Rightarrow I = \left[x \cdot \cos(\log x) + \left\{ \sin(\log x) \int 1 dx - \left(\frac{d \sin(\log x)}{dx} \int 1 \cdot dx \right) dx \right\} \right]$$

$$\Rightarrow I = \left[x \cdot \cos(\log x) + \left\{ x \cdot \sin(\log x) - \left(\cos(\log x) \frac{1}{x} x \right) dx \right\} \right]$$

$$\Rightarrow I = \left[x \cdot \cos(\log x) + \left\{ x \cdot \sin(\log x) - \left(\cos(\log x) \frac{1}{x} x \right) dx \right\} \right]$$

$$\Rightarrow I = [x \cdot \cos(\log x) + \{x \cdot \sin(\log x) - (\cos(\log x)) dx\}]$$

$$\Rightarrow I = [x \cdot \cos(\log x) + x \cdot \sin(\log x) - I]$$

$$\Rightarrow 2I = [x \cdot \cos(\log x) + x \cdot \sin(\log x)]$$

$$\Rightarrow I = \frac{x}{2} [\cos(\log x) + \sin(\log x)] + c$$

$$\text{Ans) B } \frac{x}{2} [\cos(\log x) + \sin(\log x)] + c$$

17. Question

Mark (✓) against the correct answer in each of the following:

$$\int \sec^3 x dx = ?$$

A. $\frac{1}{2} \{ \sec x \tan x - \log |\sec x + \tan x| \} + C$

B. $\frac{1}{2} \{ \sec x \tan x + \log |\sec x + \tan x| \} + C$

C. $2 \{ \sec x \tan x + \log |\sec x + \tan x| \} + C$

D. none of these

Answer**To find:** Value of $\int \sec^3 x dx$ **Formula used:**

$$(i) \int f(x)g(x)dx = f(x) \int g(x)dx - \int \left[f'(x) \int g(x)dx \right] dx$$

We have, $I = \int \sec^3 x dx \dots (i)$

$$I = \int \sec x \sec^2 x dx$$

Taking $\sec x$ as first function and $\sec^2 x$ as second function.

$$\Rightarrow I = \left[\sec x \int \sec^2 x dx - \int \left[\frac{d[\sec x]}{dx} \int \sec^2 x dx \right] dx \right]$$

$$\Rightarrow I = \left[\sec x \tan x - \int [\sec x \tan x \tan x] dx \right]$$

$$\Rightarrow I = \left[\sec x \tan x - \int [\sec x \tan^2 x] dx \right]$$

$$\Rightarrow I = \left[\sec x \tan x - \int [\sec x (\sec^2 x - 1)] dx \right]$$

$$\Rightarrow I = \left[\sec x \tan x - \int (\sec^3 x - \sec x) dx \right]$$

$$\Rightarrow I = \left[\sec x \tan x - \int \sec^3 x dx + \int \sec x dx \right]$$

$$\Rightarrow I = [\sec x \tan x - I + \log|\sec x + \tan x| + c]$$

$$\Rightarrow 2I = [\sec x \tan x + \log|\sec x + \tan x| + c]$$

$$\Rightarrow I = \frac{1}{2} [\sec x \tan x + \log|\sec x + \tan x| + c]$$

$$\text{Ans) B } \frac{1}{2} [\sec x \tan x + \log|\sec x + \tan x| + c]$$

18. Question

Mark (✓) against the correct answer in each of the following:

$$\int \left\{ \frac{1}{(\log x)} - \frac{1}{(\log x)^2} \right\} dx = ?$$

A. $x \log x + C$

B. $\frac{x}{\log x} + C$

C. $x + \frac{1}{\log x} + C$

D. none of these

Answer

To find: Value of $\int \left\{ \frac{1}{(\log x)} - \frac{1}{(\log x)^2} \right\} dx$

Formula used:

$$(i) \int f(x)g(x)dx = f(x) \int g(x)dx - \int \left[f'(x) \int g(x)dx \right] dx$$

We have, $I = \int \left\{ \frac{1}{(\log x)} - \frac{1}{(\log x)^2} \right\} dx \dots (i)$

Put $t = \log x$

$$e^t = e^{\log x} = x$$

$$\frac{dx}{dt} = e^t$$

$$\Rightarrow dx = e^t dt$$

$$\Rightarrow I = \int \left\{ \frac{1}{t} - \frac{1}{t^2} \right\} dx$$

We know $\int e^x (f(x) + f'(x)) dx = e^x f(x)$

$$\Rightarrow I = \int \left\{ \frac{1}{t} - \frac{1}{t^2} \right\} dx = e^t \frac{1}{t}$$

$$\Rightarrow \frac{x}{\log x} + c$$

Ans) B $\frac{x}{\log x} + c$



19. Question

Mark (✓) against the correct answer in each of the following:

$$\int 2x^3 e^{x^2} dx = ?$$

A. $e^{x^2} (x^2 - 1) + C$

B. $e^{x^2} (x^2 + 1) + C$

C. $e^{x^2} (x + 1) + C$

D. none of these

Answer

To find: Value of $\int \left\{ \frac{1}{(\log x)} - \frac{1}{(\log x)^2} \right\} dx$

Formula used:

$$(i) \int f(x)g(x)dx = f(x) \int g(x)dx - \int \left[f'(x) \int g(x)dx \right] dx$$

We have, $I = \int \left\{ \frac{1}{(\log x)} - \frac{1}{(\log x)^2} \right\} dx \dots (i)$

Put $t = \log x$

$$e^t = e^{\log x} = x$$

$$\frac{dx}{dt} = e^t$$

$$\Rightarrow dx = e^t dt$$

$$\Rightarrow I = \int \left\{ \frac{1}{t} - \frac{1}{t^2} \right\} dx$$

$$\text{We know } \int e^x (f(x) + f'(x)) dx = e^x f(x)$$

$$\Rightarrow I = \int \left\{ \frac{1}{t} - \frac{1}{t^2} \right\} dx = e^t \frac{1}{t}$$

$$\Rightarrow \frac{x}{\log x} + c$$

Ans) B $\frac{x}{\log x} + c$

20. Question

Mark (✓) against the correct answer in each of the following:

$$\int (x2^x) dx = ?$$

A. $\frac{2^x}{(\log 2)} (x + \log 2) + C$

B. $\frac{2^x}{(\log 2)^2} (x + \log 2 - 1) + C$

C. $\frac{x \cdot 2^x}{(\log 2)} + \frac{2^x}{(\log 2)^2} + C$

D. none of these

Answer

To find: Value of $\int (x2^x) dx$

Formula used:

$$(i) \int f(x)g(x)dx = f(x) \int g(x)dx - \int \left[f'(x) \int g(x)dx \right] dx$$

We have, $I = \int (x2^x) dx \dots (i)$

$$\Rightarrow I = x \int 2^x dx - \int \left(\frac{dx}{dx} \int 2^x dx \right) dx$$

$$\Rightarrow I = x \frac{2^x}{\log 2} - \int \left(\frac{2^x}{\log 2} \right) dx$$

$$\Rightarrow I = x \frac{2^x}{\log 2} - \frac{1}{\log 2} \int 2^x dx$$



$$\Rightarrow I = x \frac{2^x}{\log 2} - \frac{1}{\log 2} \frac{2^x}{\log 2}$$

$$\Rightarrow I = \frac{x \cdot 2^x}{\log 2} - \frac{2^x}{(\log 2)^2} + c$$

$$\Rightarrow I = \frac{2^x}{(\log 2)^2} (x \log 2 - 1) + c$$

Ans) D

21. Question

Mark (✓) against the correct answer in each of the following:

$$\int x \cot^2 x \, dx = ?$$

A. $-x \cot x + \frac{x^2}{2} + \log |\sin x| + C$

B. $-x \cot x - \frac{x^2}{2} + \log |\sin x| + C$

C. $-x \cot x + \frac{x^2}{2} - \log |\sin x| + C$

D. none of these

Answer

To find: Value of $\int x \cot^2 x \, dx$



Formula used:

$$(i) \int f(x)g(x)dx = f(x) \int g(x)dx - \int \left[f'(x) \int g(x)dx \right] dx$$

We have, $I = \int x \cot^2 x \, dx \dots (i)$

$$\Rightarrow I = x \int \cot^2 x \, dx - \int \left(\frac{dx}{dx} \int \cot^2 x \, dx \right) dx$$

$$\Rightarrow I = x \int (\operatorname{cosec}^2 x - 1) \, dx - \int \left(1 \cdot \int (\operatorname{cosec}^2 x - 1) \, dx \right) dx$$

$$\Rightarrow I = x(-\cot x - x) - \int (-\cot x - x) \, dx$$

$$\Rightarrow I = -x \cot x - x^2 + \log |\sin x| + \frac{x^2}{2}$$

$$\Rightarrow I = -x \cot x - \frac{x^2}{2} + \log |\sin x| + c$$

Ans) B $-x \cot x - \frac{x^2}{2} + \log |\sin x| + c$

22. Question

Mark (✓) against the correct answer in each of the following:

$$\int \sin \sqrt{x} \, dx = ?$$

- A. $-\sqrt{x} \cos \sqrt{x} + C$
- B. $-\sqrt{x} \cos \sqrt{x} - 2 \sin \sqrt{x} - C$
- C. $-2\sqrt{x} \cos \sqrt{x} + 2 \sin \sqrt{x} - C$
- D. none of these

Answer

To find: Value of $\int \sin \sqrt{x} dx$

Formula used: $\int \frac{1}{x} dx = \log|x| + c$

We have, $I = \int \sin \sqrt{x} dx \dots (i)$

$$\sqrt{x} = t$$

$$\Rightarrow \frac{1}{2\sqrt{x}} = \frac{dt}{dx}$$

$$\Rightarrow dx = 2\sqrt{x} dt$$

$$\Rightarrow dx = 2t dt$$

$$I = \int \sin t \cdot 2t dt$$

$$I = 2 \int t \cdot \sin t dt$$

$$\Rightarrow I = 2t \int \sin t dt - \int \left(\frac{dt}{dt} \int \sin t dt \right) dt$$

$$\Rightarrow I = 2t (-\cos t) - \int 1 (-\cos t) dt$$

$$\Rightarrow I = 2t (-\cos t) + \int \cos t dt$$

$$\Rightarrow I = 2t (-\cos t) + \sin t + c$$

$$\Rightarrow I = -2\sqrt{x} \cos \sqrt{x} + \sin \sqrt{x} + c$$

Ans) C $2\sqrt{x} \cos \sqrt{x} + \sin \sqrt{x} + c$

23. Question

Mark (✓) against the correct answer in each of the following:

$$\int e^{\sin x} \sin 2x dx = ?$$

- A. $(2 \sin x) e^{\sin x} + C$
- B. $(2 \cos x) e^{\sin x} + C$
- C. $2e^{\sin x} (\sin x + 1) + C$
- D. $2e^{\sin x} (\sin x - 1) + C$

Answer

To find: Value of $\int e^{\sin x} \sin 2x dx$

